List of Tables

	Page
Table 1.1 Telomere binding proteins and their role	47
Table 2.1 Cell seeding densities used MTT assay for cell viability and cell cycle analysis	60
Table 2.2 Composition of hypotonic fluorochrome solution used in cell cycle analysis	60
Table 2.3 Components of solutions used in the flow cytometric analysis of colorectal cell lines	66
Table 2.4 Volumes and dilutions of the antibodies used in the cell cycle analysis of colorectal cell lines	66
Table 2.5 Composition of the senescence marker solution	68
Table 3.1 Observed characteristics of each of the four colorectal cell lines chosen for initial study	73
Table 3.2 Concentration of RHPS4 causing 25%, 50% and 75% growth inhibition in a panel of four colorectal cell lines	74
Table 3.3 Concentration range of RHPS4 used to analyse the effects of senescence on HCT116, HT-29 and KM12 cells	98
Table 3.4 Comparison of the data derived from the original method used and that using equation 3.1	106
Table 6.1 How the five-compartment models was committed to COPASI	175
Table 6.2 Species involved in the five-compartment model	175
Table 6.3 The averaged rate parameters over the 21 days of the experiment according to the five-compartment model	176
Table 6.4 The average compartmental values for RHPS4 over the 21 days of the experiment according to the five-compartment model	176
Table 6.5 The average starting compartment values for doxorubicin in the five-compartment simulations using COPASI	178
Table 6.6 The average rate parameter values for each concentration of doxorubicin over the 21 day period according to the five-compartment mathematical model as applied to COPASI	178
Table 6.7 Rate parameters calculated when applying 0.1x, 0.15x and 0.3x compartment data to the five-compartment mathematical model	183
Table 6.8 The original data from doxorubicin and the data used in COPASI following parameter fitting	190
Table 6.9 The original data from RHPS4 when applied to the five-compartment mathematical model and when used in COPASI following parameter fitting	192